



# Production of glycolipid biosurfactants, mannosylerythritol lipids, from pentoses and D-glucose/D-xylose mixtures by *Pseudozyma* yeast strains



Nuno Torres Faria<sup>a,b,c</sup>, Marisa V. Santos<sup>a</sup>, Pedro Fernandes<sup>a,d</sup>, Luís Lopes Fonseca<sup>e,1</sup>, César Fonseca<sup>c,\*</sup>, Frederico Castelo Ferreira<sup>a,b,\*\*</sup>

<sup>a</sup> Department of Bioengineering and IBB-Institute for Biotechnology and Bioengineering, Instituto Superior Técnico, Universidade de Lisboa, Av. Rovisco Pais, 1049-001 Lisboa, Portugal

<sup>b</sup> MIT-Portugal Program, 77 Massachusetts Avenue, E40-221, Cambridge, MA 02139, USA

<sup>c</sup> Laboratório Nacional de Energia e Geologia, I.P., Unidade de Bioenergia, Estrada do Paço do Lumiar 22, 1649-038 Lisboa, Portugal

<sup>d</sup> Faculdade de Engenharia, Universidade Lusófona de Humanidades e Tecnologias, Av. Campo Grande 376, 1749-024 Lisboa, Portugal

<sup>e</sup> Instituto de Tecnologia Química e Biológica (ITQB), Universidade Nova de Lisboa, 2780-156 Oeiras, Portugal

## ARTICLE INFO

### Article history:

Received 24 June 2014

Received in revised form 28 July 2014

Accepted 14 August 2014

Available online 23 August 2014

### Keywords:

Mannosylerythritol lipids (MEL)

Biosurfactants

Pentoses

D-Xylose

*Pseudozyma* spp.

## ABSTRACT

The pentose-assimilating capacity of yeasts from the genus *Pseudozyma*, *P. antarctica* PYCC 5048<sup>T</sup>, *P. aphidis* PYCC 5535<sup>T</sup> and *P. rugulosa* PYCC 5537<sup>T</sup>, was exploited towards the production of mannosylerythritol lipids (MEL), a glycolipid with biosurfactant properties. The three strains tested were able to grow on D-xylose and L-arabinose with similar maximum specific growth rates to those estimated on D-glucose (around 0.2 h<sup>-1</sup>). The highest MEL titres (4.8–5.4 g/l) and yields (0.11–0.14 g/g) from D-xylose were found in *P. antarctica* PYCC 5048<sup>T</sup>, which presented similar values to those estimated on D-glucose and on D-xylose/D-glucose mixtures. *P. rugulosa* PYCC 5537<sup>T</sup> showed a pattern of sugar conversion into MEL similar to *P. antarctica*, but at 40% lower titres. *P. aphidis* PYCC 5535<sup>T</sup> presented lower MEL titres from D-xylose (1.2 g/l) than from D-glucose (3.4 g/l). Nitrate supply increased sugar consumption rate and, when accompanied by D-glucose or D-xylose feeding, additional biomass production. In this case, sugar was completely consumed before sugar feeding at day 7, but not when feeding is performed at day 4. Higher MEL titres were obtained for the later condition reaching 7.3 g/l and 5.8 g/l, in fed-batch cultures with glucose and xylose, respectively.

© 2014 Elsevier Ltd. All rights reserved.

## 1. Introduction

Microbial surfactants are recognized as fine chemicals with high-value applications and interesting characteristics such as low toxicity, high biodegradability, effectiveness at extreme temperatures or pH, and mild production conditions when compared to

chemical surfactants [1–3]. Biosurfactants, in particular lipopeptides and glycolipids, have industrial applications in the production of food, cosmetics, and pharmaceuticals, as well as for soil decontamination of heavy metals, oils and other toxic organics [4]. In microbial environments, the roles of biosurfactants include the increase of surface area and bioavailability of hydrophobic water-insoluble substrates, heavy metal binding, bacterial pathogenesis, quorum sensing and biofilm formation [2].

Mannosylerythritol lipids (MELs) are glycolipids containing a 4-O-β-D-mannopyranosyl-meso-erythritol as the glycosidic/hydrophilic moiety and two short-chain fatty acids (usually C8–C12) as the hydrophobic groups (Fig. 1) [4]. These amphiphilic properties classify these molecules as biosurfactants [1]. MELs are designated as MEL-A, -B, -C, according with their elution position in thin layer chromatography (TLC), and corresponding to the degree of acetylation at C-4 and/or at C-6 position of the mannosyl moiety [5]. MEL-A corresponds to the diacetylated compound while MEL-B and MEL-C are monoacetylated at C-6 and C-4 of

\* Corresponding author at: Laboratório Nacional de Energia e Geologia, I.P., Unidade de Bioenergia, Estrada do Paço do Lumiar 22, 1649-038 Lisboa, Portugal. Tel.: +351 210924717.

\*\* Corresponding author at: Department of Bioengineering and IBB-Institute for Biotechnology and Bioengineering, Instituto Superior Técnico, Universidade de Lisboa, Av. Rovisco Pais, 1049-001 Lisboa, Portugal. Tel.: +351 218419598.

E-mail addresses: [cesar.fonseca@lneg.pt](mailto:cesar.fonseca@lneg.pt) (C. Fonseca), [frederico.ferreira@ist.utl.pt](mailto:frederico.ferreira@ist.utl.pt), [frederico.castelo@yahoo.com](mailto:frederico.castelo@yahoo.com) (F.C. Ferreira).

<sup>1</sup> Present address: Integrative BioSystems Institute and The Wallace H. Coulter Department of Biomedical Engineering, Georgia Institute of Technology, 313 Ferst Drive, Atlanta, GA 30332, USA.